

Wiener-Hopf plus Hankel operators on the real line with unitary and sectorial symbols

G. Bogveradze and L. P. Castro

ABSTRACT. Wiener-Hopf plus Hankel operators acting between Lebesgue spaces on the real line are studied in view of their invertibility, one sided-invertibility, Fredholm, and semi-Fredholm properties. This is done in two different cases: (i) when the Fourier symbols of the operators are unitary functions, and (ii) when the Fourier symbols are related with sectorial elements appearing in factorizations of functions originated by the Fourier symbols of the operators. The obtained result for the case (i) may be viewed as a Douglas-Sarason type theorem for Wiener-Hopf plus Hankel operators.

1. Introduction

Algebraic sums of Wiener-Hopf and Hankel operators have been receiving increasing attention in the last years (cf. [1], [5], [6], [10], [12]–[19], [22], [23]). A great part of the interest is directly originated by concrete mathematical-physics applications where this kind of operators appear. This is the case of problems in wave diffraction phenomena which are modelled by boundary-transmission value problems that can be equivalently translated into systems of integral equations characterized by such kind of operators (see, e.g., [7]–[9]).

It is therefore easy to understand that a fundamental goal in those studies consists in obtaining conditions that lead to the *regularity properties* of the operators (i.e., their invertibility, Fredholm property, and other properties that are directly dependent on the kernel and image of the operators). For some classes of the so-called *Fourier symbols of the operators* their regularity properties are already known. This is the case of the continuous or, even, piecewise continuous functions on the compactified real line (cf. the above references). Despite those advances, for some other classes of Fourier symbols a complete description of the regularity properties is still missing.

In the present paper, starting simply with essential bounded Fourier symbols, we look for conditions that may ensure the above pointed properties for Wiener-Hopf plus Hankel operators having that Fourier symbols. All the detailed definitions are given in the next section.

2000 *Mathematics Subject Classification.* 47B35, 47A68, 47A53.

Key words and phrases. Wiener-Hopf plus Hankel operator, invertibility, Fredholm property, unitary function, sectorial function.

G. Bogveradze is sponsored by a grant from the Research Unit *Mathematics and Applications* through *Fundação para a Ciência e a Tecnologia* (Portugal).

2. The Wiener-Hopf plus Hankel operators in study

We will consider Wiener-Hopf plus Hankel operators of the form

$$(2.1) \quad WH_\varphi = W_\varphi + H_\varphi : L_+^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}_+) ,$$

with W_φ and H_φ being Wiener-Hopf and Hankel operators defined by

$$(2.2) \quad W_\varphi = r_+ \mathcal{F}^{-1} \varphi \cdot \mathcal{F} : L_+^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}_+) ,$$

$$(2.3) \quad H_\varphi = r_+ \mathcal{F}^{-1} \varphi \cdot \mathcal{F} J : L_+^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}_+) ,$$

respectively. As usual, $L^2(\mathbb{R})$ and $L^2(\mathbb{R}_+)$ denote the Hilbert spaces of complex-valued Lebesgue measurable functions φ , for which $|\varphi|^2$ is integrable on $\mathbb{R}$ and $\mathbb{R}_+$, respectively. In (2.1)–(2.3), $L_+^2(\mathbb{R})$ denotes the subspace of $L^2(\mathbb{R})$ formed by all functions supported in the closure of $\mathbb{R}_+ = (0, +\infty)$, the operator r_+ performs the restriction from $L^2(\mathbb{R})$ into $L^2(\mathbb{R}_+)$, $\mathcal{F}$ denotes the Fourier transformation, J is the reflection operator given by the rule $J\varphi(x) = \tilde{\varphi}(x) = \varphi(-x)$, $x \in \mathbb{R}$, and $\varphi \in L^\infty(\mathbb{R})$ is the so-called Fourier symbol.

Let $\mathbb{C}_- = \{z \in \mathbb{C} : \Im z < 0\}$ and $\mathbb{C}_+ = \{z \in \mathbb{C} : \Im z > 0\}$. As usual, let us denote by $H^\infty(\mathbb{C}_\pm)$ the set of all bounded and analytic functions in $\mathbb{C}_\pm$. *Fatou's Theorem* ensures that functions in $H^\infty(\mathbb{C}_\pm)$ have non-tangential limits on $\mathbb{R}$ almost everywhere. Thus, let $H_\pm^\infty(\mathbb{R})$ be the set of all functions in $L^\infty(\mathbb{R})$ that are non-tangential limits of elements in $H^\infty(\mathbb{C}_\pm)$. Moreover, it is well known that $H_\pm^\infty(\mathbb{R})$ are closed subalgebras of $L^\infty(\mathbb{R})$.

We will concentrate on obtaining regularity properties of this kind of Wiener-Hopf plus Hankel operators (2.1) in the case where the Fourier symbol φ is an unitary-valued function or a function somehow associated to a sectorial element through a certain factorization (and in these two cases we will make use of the just presented $H_\pm^\infty := H_\pm^\infty(\mathbb{R})$ spaces). We recall that $\phi \in L^\infty(\mathbb{R})$ is called *unitary* if $|\phi| = 1$, and is called *sectorial* if the essential range of ϕ is contained in an open half-plane whose boundary passes through the origin.

As mentioned, within the context of the present paper we are using the name of *regularity properties* of a linear operator acting between Hilbert spaces for those proprieties that arise from a direct influence of the kernel and the image of those operators. In more detail, let $T : X \rightarrow Y$ be a bounded linear operator acting between Hilbert spaces and, if $\text{Im } T$ is closed (i.e. T is *normally solvable*), let us consider the cokernel of T to be defined by the quotient $\text{Coker } T = Y/\text{Im } T$. Then, a normally solvable operator T is said to be *right-Fredholm* if $\dim \text{Coker } T$ is finite, *left-Fredholm* if $\dim \text{Ker } T$ is finite, and *Fredholm* if both $\dim \text{Ker } T$ and $\dim \text{Coker } T$ are finite. Additionally, we say that T is *left-invertible* or *right-invertible* if there exist $T_l^- : Y \rightarrow X$ or $T_r^- : Y \rightarrow X$ such that $T_l^- T = I_X$ or $TT_r^- = I_Y$, respectively. As usual, in the case when both T_l^- and T_r^- exist and additionally $T_l^- = T_r^-$, the operator T is said to be *invertible* (or *both sided-invertible*). Alternatively, it can be shown that T is left-invertible if and only if T is injective and normally solvable. In the same way, T is right-invertible if and only if T is normally solvable and surjective.

3. A Douglas–Sarason type theorem

For a Banach algebra B , we are going to denote by $\mathcal{G}B$ the group of all invertible elements in B .

We point out that if $\phi \in \mathcal{GL}^\infty(\mathbb{R})$, then there is a function $h \in \mathcal{GH}_+^\infty(\mathbb{R})$ such that $|\phi| = |h|$ almost everywhere in $\mathbb{R}$. This is only a particular case of the *Pousson–Rabindranathan Theorem* [20, 21] (cf. Theorem 4.3 below) but give a hint to the strategy behind the next results where distances between certain classes and the Fourier symbols of the operators are considered.

In the Wiener-Hopf operators case there is a well known theorem – due to Douglas and Sarason – about the Fredholm theory of this kind of operators and the distances between the Fourier symbol to certain spaces. More precisely, the theorem may be written in the following form.

THEOREM 3.1 (Douglas and Sarason [11]). *If $\varphi \in L^\infty(\mathbb{R})$ is unitary, then:*

- (a) W_φ is invertible if and only if $\text{dist}(\varphi, \mathcal{GH}_+^\infty) < 1$ or if and only if $\text{dist}(\varphi, \mathcal{GH}_-^\infty) < 1$;
- (b) W_φ is left-invertible if and only if $\text{dist}(\varphi, H_+^\infty) < 1$;
- (b') W_φ is right-invertible if and only if $\text{dist}(\varphi, H_-^\infty) < 1$;
- (c) W_φ is Fredholm if and only if $\text{dist}(\varphi, \mathcal{G}[C(\dot{\mathbb{R}}) + H_+^\infty]) < 1$ or if and only if $\text{dist}(\varphi, \mathcal{G}[C(\dot{\mathbb{R}}) + H_-^\infty]) < 1$;
- (d) W_φ is left-Fredholm if and only if $\text{dist}(\varphi, C(\dot{\mathbb{R}}) + H_+^\infty) < 1$;
- (d') W_φ is right-Fredholm if and only if $\text{dist}(\varphi, C(\dot{\mathbb{R}}) + H_-^\infty) < 1$.

As usual, $C(\dot{\mathbb{R}})$ denotes the space of all bounded continuous (complex-valued) functions on $\mathbb{R}$ for which both limits at $\pm\infty$ exist and coincide.

The last theorem was the starting motivation for obtaining such kind of result for our Wiener-Hopf plus Hankel operators. It is clear that adding an Hankel operator to the above Wiener-Hopf operator will influence several changes in the regularity properties of the resulting operator.

To consider such situation, we will use the notion of Δ -relation after extension introduced in [4] for bounded linear operators acting between Hilbert spaces, e.g. $T : X_1 \rightarrow X_2$ and $S : Y_1 \rightarrow Y_2$. We say that T is Δ -related after extension to S (and use the abbreviation $T \overset{*}{\Delta} S$) if there is an auxiliary bounded linear operator acting between Hilbert spaces $T_\Delta : X_{1\Delta} \rightarrow X_{2\Delta}$, and bounded invertible operators E and F such that

$$(3.1) \quad \begin{bmatrix} T & 0 \\ 0 & T_\Delta \end{bmatrix} = E \begin{bmatrix} S & 0 \\ 0 & I_Z \end{bmatrix} F,$$

where Z is an additional Hilbert space and I_Z represents the identity operator in Z . In the particular case where $T_\Delta = I_{X_{1\Delta}} : X_{1\Delta} \rightarrow X_{2\Delta} = X_{1\Delta}$ is the identity operator, we say the T and S are *equivalent after extension operators*. In the simplest case of $T = E S F$ (for some boundedly invertible operators E and F) we say that T and S are *equivalent operators*.

THEOREM 3.2. *Let φ be unitary in $L^\infty(\mathbb{R})$.*

- (a) WH_φ and $W_\varphi - H_\varphi$ are invertible if and only if

$$\text{dist}(\widetilde{\varphi\varphi^{-1}}, \mathcal{GH}_+^\infty) < 1$$

$$\text{or if and only if } \text{dist}(\widetilde{\varphi\varphi^{-1}}, \mathcal{GH}_-^\infty) < 1.$$

(b) WH_φ and $W_\varphi - H_\varphi$ are left-invertible if and only if

$$\text{dist}(\widetilde{\varphi\varphi^{-1}}, H_+^\infty) < 1.$$

(b') WH_φ and $W_\varphi - H_\varphi$ are right-invertible if and only if

$$\text{dist}(\widetilde{\varphi\varphi^{-1}}, H_-^\infty) < 1.$$

(c) WH_φ and $W_\varphi - H_\varphi$ are Fredholm if and only if

$$\text{dist}(\widetilde{\varphi\varphi^{-1}}, \mathcal{G}[C(\dot{\mathbb{R}}) + H_+^\infty]) < 1,$$

or if and only if $\text{dist}(\widetilde{\varphi\varphi^{-1}}, \mathcal{G}[C(\dot{\mathbb{R}}) + H_-^\infty]) < 1.$

(d) WH_φ and $W_\varphi - H_\varphi$ are left-Fredholm if and only if

$$\text{dist}(\widetilde{\varphi\varphi^{-1}}, C(\dot{\mathbb{R}}) + H_+^\infty) < 1.$$

(d') WH_φ and $W_\varphi - H_\varphi$ are right-Fredholm if and only if

$$\text{dist}(\widetilde{\varphi\varphi^{-1}}, C(\dot{\mathbb{R}}) + H_-^\infty) < 1.$$

PROOF. Our proof is based on the notion of Δ -relation after extension, which was recalled above for bounded linear operators acting between Hilbert spaces. From [4, Example 1.7] we can derive that $T = WH_\varphi$ is Δ -related after extension to the Wiener-Hopf operator $S = W_{\varphi\varphi^{-1}}$, cf. also [18, Lemma 2.1] or [19, Lemma 1] where this relation is given in an explicit way for $T_\Delta = W_\varphi - H_\varphi$ in (3.1) (or for an operator T_Δ equivalent to $W_\varphi - H_\varphi$). Thus, we are now going to analyze the Fourier symbol $\varphi\varphi^{-1}$.

Let φ^* stands for the complex conjugate of φ . Since φ is unitary, then $\varphi\varphi^* = \varphi^*\varphi = 1$, and whence $\varphi^{-1} = \varphi^*$. From here we also obtain $\widetilde{\varphi}^{-1} = \widetilde{\varphi}^*$. Observing that

$$[\widetilde{\varphi\varphi^*}] = [\widetilde{\varphi^*\varphi}] = 1,$$

we reach to $\widetilde{\varphi}\widetilde{\varphi}^* = \widetilde{\varphi}^*\widetilde{\varphi} = 1$. This means that $\widetilde{\varphi}$ is unitary and so it is $\varphi\varphi^{-1}$. We can now apply Theorem 3.1 to the operator $W_{\varphi\varphi^{-1}}$ and obtain all the above stated conditions in terms of distances. Now, the result follows if we interpret (3.1) as an equivalence after extension relation between $\text{diag}[T, T_\Delta] = \text{diag}[WH_\varphi, W_\varphi - H_\varphi]$ and $S = W_{\varphi\varphi^{-1}}$. $\square$

REMARK 3.3. We would like to point out that in the above reasoning (of the Δ -relation after extension) it is not possible to exclude the factor $\widetilde{\varphi}^{-1}$ for obtaining a “pure” (direct) result depending only on φ (without extra factors). Consider for instance the case (b). From the condition $\text{dist}(\widetilde{\varphi\varphi^{-1}}, H_+^\infty) < 1$ it follows that

$$\inf_{h \in H_+^\infty} \|\widetilde{\varphi\varphi^{-1}} - h\|_{L^\infty(\mathbb{R})} < 1,$$

which is equivalent to

$$\inf_{h \in H_+^\infty} \|\varphi - h\widetilde{\varphi}\|_{L^\infty(\mathbb{R})} < 1,$$

but this is the distance from φ to the space $\{h\widetilde{\varphi} : h \in H_+^\infty \text{ and } \varphi \text{ is unitary}\}$. Note that this space is obviously different from H_+^∞ . In fact, if we take $\varphi \in H_+^\infty$, with $\varphi(x) = e^{i\lambda x}$ and $\lambda > 0$ ($x \in \mathbb{R}$), then choosing $h(x) = e^{i\mu x}$ with $0 < \mu < \lambda$ we have $h\widetilde{\varphi} \in H_-^\infty$. This also means that the space is directly dependent on the structure of $\widetilde{\varphi}$ (and therefore on $\widetilde{\varphi}^{-1}$).

4. Sectorial Symbols

In the present section we will work with the so-called *sectorial Fourier symbols*.

DEFINITION 4.1. A function $f \in L^\infty(\mathbb{R})$ is said to be *sectorial* if there exist a real number $\varepsilon > 0$ and a $c \in \mathbb{C}$ of modulus 1 such that

$$\Re(cf(x)) \geq \varepsilon,$$

almost everywhere on $\mathbb{R}$.

We will denote by $\mathcal{S}$ the set of all sectorial functions (in $L^\infty(\mathbb{R})$). Once again, for Wiener-Hopf operators with such kind of Fourier symbols a description of the possible regularity properties is known. Additionally, more detailed information (upon some factorizations of the original Fourier symbols) is also known in the following form.

THEOREM 4.2. [2, Theorem 6.17] *If $\varphi \in \mathcal{GL}^\infty(\mathbb{R})$, then:*

- (a) W_φ is invertible if and only if $\varphi = sh$, $s \in \mathcal{S}$, $h \in \mathcal{GH}_\pm^\infty$;
- (b) W_φ is left-invertible if and only if $\varphi = sh$, $s \in \mathcal{S}$, $h \in H_+^\infty$;
- (b') W_φ is right-invertible if and only if $\varphi = sh$, $s \in \mathcal{S}$, $h \in H_-^\infty$;
- (c) W_φ is Fredholm if and only if $\varphi = sh$, $s \in \mathcal{S}$, $h \in \mathcal{G}[C(\dot{\mathbb{R}}) + H_\pm^\infty]$;
- (d) W_φ is left-Fredholm if and only if $\varphi = sh$, $s \in \mathcal{S}$, $h \in C(\dot{\mathbb{R}}) + H_+^\infty$;
- (d') W_φ is right-Fredholm if and only if $\varphi = sh$, $s \in \mathcal{S}$, $h \in C(\dot{\mathbb{R}}) + H_-^\infty$.

Such type of theorem is also valid for Wiener-Hopf plus Hankel operators, involving naturally some modifications over the symbol conditions. To prove the next theorem in this direction, we first need to recall some auxiliary results.

THEOREM 4.3 (Pousson and Rabindranathan [20, 21]). *If $a \in \mathcal{GL}^\infty(\mathbb{R})$, then there exist a unitary-valued u and $h \in \mathcal{GH}_+^\infty$ such that $a = uh$, almost everywhere on $\mathbb{R}$.*

LEMMA 4.4. [3, Lemma 2.21] *If E is a subset of $L^\infty(\mathbb{R})$, $\varphi \in L^\infty(\mathbb{R})$ is unitary-valued, and $\text{dist}(\varphi, E) < 1$, then there exist a function $f \in E$ and a sectorial function $s \in \mathcal{GL}^\infty(\mathbb{R})$ such that $\varphi = sf$.*

Now we introduce and prove a Wiener-Hopf plus/minus Hankel version of the Theorem 4.2.

THEOREM 4.5. *Let $\varphi \in \mathcal{GL}^\infty(\mathbb{R})$.*

- (a) WH_φ and $W_\varphi - H_\varphi$ are both invertible if and only if

$$\widetilde{\varphi\varphi^{-1}} = sh, \quad s \in \mathcal{S}, \quad h \in \mathcal{GH}_\pm^\infty.$$

- (b) WH_φ and $W_\varphi - H_\varphi$ are both left-invertible if and only if

$$\widetilde{\varphi\varphi^{-1}} = sh, \quad s \in \mathcal{S}, \quad h \in H_+^\infty.$$

- (c) WH_φ and $W_\varphi - H_\varphi$ are both right-invertible if and only if

$$\widetilde{\varphi\varphi^{-1}} = sh, \quad s \in \mathcal{S}, \quad h \in H_-^\infty.$$

- (d) WH_φ and $W_\varphi - H_\varphi$ are both Fredholm if and only if

$$\widetilde{\varphi\varphi^{-1}} = sh_1h_2, \quad s \in \mathcal{S}, \quad h_1 \in \mathcal{G}[C(\dot{\mathbb{R}}) + H_\pm^\infty], \quad h_2 \in \mathcal{GH}_\pm^\infty.$$

(e) WH_φ and $W_\varphi - H_\varphi$ are both left-Fredholm if and only if

$$\widetilde{\varphi\varphi^{-1}} = sh_1h_2, \quad s \in \mathcal{S}, \quad h_1 \in C(\dot{\mathbb{R}}) + H_+^\infty, \quad h_2 \in \mathcal{G}H_+^\infty.$$

(f) WH_φ and $W_\varphi - H_\varphi$ are both right-Fredholm if and only if

$$\widetilde{\varphi\varphi^{-1}} = sh_1h_2, \quad s \in \mathcal{S}, \quad h_1 \in C(\dot{\mathbb{R}}) + H_-^\infty, \quad h_2 \in \mathcal{G}H_-^\infty.$$

PROOF. First we will prove the “+”-version (cf. the indices). As a global property, from the Δ -relation after extension we know that $WH_\varphi \overset{*}{\Delta} W_{\varphi\varphi^{-1}}$. Additionally, from the Theorem 4.3 (of Pousson and Rabindranathan), we have

$$(4.1) \quad \widetilde{\varphi\varphi^{-1}} = u_\varphi h_\varphi,$$

where u_φ is unitary-valued, and $h_\varphi \in \mathcal{G}H_+^\infty$ (the notation of φ in index is only to reinforce that those factors depend on φ). Therefore, (4.1) yields that $W_{\varphi\varphi^{-1}}$ is equivalent to W_{u_φ} . Using now the Douglas–Sarason Theorem (cf. Theorem 3.1), we have:

- (i) W_{u_φ} is invertible if and only if $\text{dist}(u_\varphi, \mathcal{G}H_+^\infty) < 1$;
- (ii) W_{u_φ} is left-invertible if and only if $\text{dist}(u_\varphi, H_+^\infty) < 1$;
- (iii) W_{u_φ} is Fredholm if and only if $\text{dist}(u_\varphi, \mathcal{G}[C(\dot{\mathbb{R}}) + H_+^\infty]) < 1$;
- (iv) W_{u_φ} is left-Fredholm if and only if $\text{dist}(u_\varphi, C(\dot{\mathbb{R}}) + H_+^\infty) < 1$.

This combined with the above Lemma 4.4 yields that:

- (i') If W_{u_φ} is invertible, then $u_\varphi = sh$ for some $s \in \mathcal{S}$ and $h \in \mathcal{G}H_+^\infty$;
- (ii') If W_{u_φ} is left-invertible, then $u_\varphi = sh$ for some $s \in \mathcal{S}$ and $h \in H_+^\infty$;
- (iii') If W_{u_φ} is Fredholm, then $u_\varphi = sh$ for some $s \in \mathcal{S}$ and $h \in \mathcal{G}[C(\dot{\mathbb{R}}) + H_+^\infty]$;
- (iv') If W_{u_φ} is left-Fredholm, then $u_\varphi = sh$ for some $s \in \mathcal{S}$ and $h \in C(\dot{\mathbb{R}}) + H_+^\infty$.

Using the multiplication by h_φ in the last four identities (and recalling the identity $\widetilde{\varphi\varphi^{-1}} = u_\varphi h_\varphi$), we conclude that:

- (i'') If W_{u_φ} is invertible, then

$$\widetilde{\varphi\varphi^{-1}} = sh, \quad s \in \mathcal{S}, \quad h \in \mathcal{G}H_+^\infty;$$

- (ii'') If W_{u_φ} is left-invertible, then

$$\widetilde{\varphi\varphi^{-1}} = sh, \quad s \in \mathcal{S}, \quad h \in H_+^\infty;$$

- (iii'') If W_{u_φ} is Fredholm, then

$$\widetilde{\varphi\varphi^{-1}} = sh_1h_2, \quad s \in \mathcal{S}, \quad h_1 \in \mathcal{G}[C(\dot{\mathbb{R}}) + H_+^\infty], \quad h_2 \in \mathcal{G}H_+^\infty;$$

- (iv'') If W_{u_φ} is left-Fredholm, then

$$\widetilde{\varphi\varphi^{-1}} = sh_1h_2, \quad s \in \mathcal{S}, \quad h_1 \in C(\dot{\mathbb{R}}) + H_+^\infty, \quad h_2 \in \mathcal{G}H_+^\infty$$

where h_1 and h_2 represent h and h_φ , respectively.

Now we will proceed with the reverse implications. Let us then assume the following four situations:

- (j) $\widetilde{\varphi\varphi^{-1}} = sh$, for some $s \in \mathcal{S}$ and $h \in \mathcal{G}H_+^\infty$;
- (jj) $\widetilde{\varphi\varphi^{-1}} = sh$, for some $s \in \mathcal{S}$ and $h \in H_+^\infty$;

- (jjj) $\widetilde{\varphi\varphi^{-1}} = sh_1h_2$, for some $s \in \mathcal{S}$, $h_1 \in \mathcal{G}[C(\dot{\mathbb{R}}) + H_+^\infty]$ and $h_2 \in \mathcal{GH}_+^\infty$;
- (jv) $\widetilde{\varphi\varphi^{-1}} = sh_1h_2$, for some $s \in \mathcal{S}$, $h_1 \in C(\dot{\mathbb{R}}) + H_+^\infty$ and $h_2 \in \mathcal{GH}_+^\infty$.

In the first two cases (j) and (jj) we directly obtain from Theorem 4.2 that $\widetilde{W_{\varphi\varphi^{-1}}}$ is invertible or left-invertible, respectively. Therefore, the above mentioned Δ -relation after extension yields the same properties for the diagonal matrix operator $\text{diag}[WH_\varphi, W_\varphi - H_\varphi]$, and the result follows (for these two cases).

In the cases (jjj) and (jv), formula (4.1) yields that

$$W_{u_\varphi} = W_s W_{h_1} W_{h_2 h_\varphi^{-1}}.$$

Note that W_s is an invertible operator because s is sectorial, and $W_{h_2 h_\varphi^{-1}}$ is also invertible due to $h_2 h_\varphi^{-1} \in \mathcal{GH}_+^\infty$ (having by inverse operator the corresponding Wiener-Hopf operator with Fourier symbol $h_2^{-1} h_\varphi$). Therefore, W_{u_φ} is equivalent to W_{h_1} . Thus, to conclude the desired result by the use of the Δ -relation after extension, we only need to describe the analysis of W_{h_1} in those corresponding cases:

- (k) If $h_1 \in \mathcal{G}[C(\dot{\mathbb{R}}) + H_+^\infty]$, then W_{h_1} is Fredholm;
- (kk) If $h_1 \in C(\dot{\mathbb{R}}) + H_+^\infty$, then W_{h_1} is left-Fredholm.

In fact, propositions (k) and (kk) arise from the *Douglas Theorem* (cf., e.g., [2, Theorem 5.7]), and by observing that $h_1 \in \mathcal{GL}^\infty(\mathbb{R})$ (because $\widetilde{\varphi\varphi^{-1}} \in \mathcal{GL}^\infty(\mathbb{R})$, $h_2 \in \mathcal{GH}_+^\infty$ and $s \in \mathcal{GL}^\infty(\mathbb{R})$).

We have therefore proved the “+”-version of the theorem. Now, by passage to adjoints, the remaining “-”-versions are obtained. $\square$

The “only if” part of Theorem 4.5, directly yields the main conclusion of the present work:

COROLLARY 4.6. Let $\varphi \in \mathcal{GL}^\infty(\mathbb{R})$.

- (a) If $\widetilde{\varphi\varphi^{-1}} = sh$, with $s \in \mathcal{S}$ and $h \in \mathcal{GH}_\pm^\infty$, then WH_φ is invertible.
- (b) If $\widetilde{\varphi\varphi^{-1}} = sh$, with $s \in \mathcal{S}$ and $h \in H_+^\infty$, then WH_φ is left-invertible.
- (c) If $\widetilde{\varphi\varphi^{-1}} = sh$, with $s \in \mathcal{S}$ and $h \in H_-^\infty$, then WH_φ is right-invertible.
- (d) If $\widetilde{\varphi\varphi^{-1}} = sh_1h_2$, with $s \in \mathcal{S}$, $h_1 \in \mathcal{G}[C(\dot{\mathbb{R}}) + H_\pm^\infty]$, and $h_2 \in \mathcal{GH}_\pm^\infty$, then WH_φ is Fredholm.
- (e) If $\widetilde{\varphi\varphi^{-1}} = sh_1h_2$, with $s \in \mathcal{S}$, $h_1 \in C(\dot{\mathbb{R}}) + H_+^\infty$ and $h_2 \in \mathcal{GH}_+^\infty$, then WH_φ is left-Fredholm.
- (f) If $\widetilde{\varphi\varphi^{-1}} = sh_1h_2$, with $s \in \mathcal{S}$, $h_1 \in C(\dot{\mathbb{R}}) + H_-^\infty$ and $h_2 \in \mathcal{GH}_-^\infty$, then WH_φ is right-Fredholm.

We end with an example showing the applicability of the last result. Let us consider the Wiener-Hopf plus Hankel operator

$$WH_{\varphi_p} : L_+^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}_+),$$

with the particular Fourier symbol

$$\varphi_p(x) = (2 + \sin x) e^{i\alpha x}, \quad x \in \mathbb{R},$$

where $\alpha \in \mathbb{R}$ is a given parameter. Direct computations show that

$$\varphi_p(x) \widetilde{\varphi_p^{-1}}(x) = \frac{2 + \sin x}{2 - \sin x} e^{2i\alpha x}.$$

So, if we choose $s_p(x) = (2 + \sin x)/(2 - \sin x)$ and $h_p(x) = e^{2i\alpha x}$, we see that $s_p \in \mathcal{S}$. This occurs because $1/3 \leq s_p \leq 3$, and therefore the range of s_p is contained in the right half-plane (which boundary passes through the origin). Moreover, depending whether $\alpha \geq 0$ or $\alpha \leq 0$, we have $h_p \in H_+^\infty$ or $h_p \in H_-^\infty$, respectively. Therefore, applying Corollary 4.6 to

$$\varphi_p \widetilde{\varphi_p^{-1}} = s_p h_p ,$$

we conclude that:

- (a) if $\alpha = 0$, then WH_{φ_p} is invertible;
- (b) if $\alpha > 0$, then WH_{φ_p} is left-invertible;
- (c) if $\alpha < 0$, then WH_{φ_p} is right-invertible.

Acknowledgements: The work was supported by *Fundação para a Ciência e a Tecnologia* through *Unidade de Investigação Matemática e Aplicações* of University of Aveiro, Portugal.

References

- [1] E.L. Basor and T. Ehrhardt, Factorization theory for a class of Toeplitz + Hankel operators, *J. Oper. Theory*, **51**(2004), 411–433.
- [2] A. Böttcher, Yu.I. Karlovich, I.M. Spitkovsky, *Convolution Operators and Factorization of Almost Periodic Matrix Functions*, Birkhäuser Verlag, Basel (2002).
- [3] A. Böttcher, B. Silbermann, *Analysis of Toeplitz Operators*, Springer-Verlag, Berlin (1990).
- [4] L.P. Castro and F.-O. Speck, Regularity properties and generalized inverses of delta-related operators, *Z. Anal. Anwendungen*, **17** (1998), 577–598.
- [5] L.P. Castro and F.-O. Speck, Inversion of matrix convolution type operators with symmetry, *Port. Math. (N.S.)*, **62** (2005), 193–216.
- [6] L.P. Castro, F.-O. Speck and F.S. Teixeira, A direct approach to convolution type operators with symmetry, *Math. Nachr.*, **269/270** (2004), 73–85.
- [7] L.P. Castro, F.-O. Speck and F.S. Teixeira, Explicit solution of a Dirichlet-Neumann wedge diffraction problem with a strip, *J. Integral Equations Appl.*, **15** (2003), 359–383.
- [8] L.P. Castro, F.-O. Speck and F.S. Teixeira, On a class of wedge diffraction problems posted by Erhard Meister, *Oper. Theory Adv. Appl.*, **147** (2004), 213–240.
- [9] L.P. Castro, F.-O. Speck and F.S. Teixeira, Mixed boundary value problems for the Helmholtz equation in a quadrant, *Integral Equations Operator Theory*, 44 pp., in press.
- [10] A.C. Conceição, V.G. Kravchenko and F.S. Teixeira, Factorization of some classes of matrix functions and the resolvent of a Hankel operator, in: *Factorization, Singular Operators and Related Problems*, Kluwer Acad. Publ., Dordrecht (2003), pp. 101–110.
- [11] R.G. Douglas and D. Sarason, Fredholm Toeplitz operators, *Proc. Amer. Math. Soc.*, **26** (1970), 117–120.
- [12] T. Ehrhardt, Invertibility theory for Toeplitz plus Hankel operators and singular integral operators with flip, *J. Funct. Anal.*, **208** (2004), 64–106.
- [13] N. Karapetians and S. Samko, *Equations with Involution Operators*, Birkhäuser, Boston (2001).
- [14] V.G. Kravchenko, A.B. Lebre, G.S. Litvinchuk and F.S. Teixeira, Fredholm theory for a class of singular integral operators with Carleman shift and unbounded coefficients, *Math. Nachr.*, **172** (1995), 199–210.
- [15] V.G. Kravchenko and G.S. Litvinchuk, *Introduction to the Theory of Singular Integral Operators with Shift*, Kluwer Academic Publishers Group, Dordrecht (1994).
- [16] A.B. Lebre, E. Meister and F.S. Teixeira, Some results on the invertibility of Wiener-Hopf-Hankel operators, *Z. Anal. Anwendungen*, **11** (1992), 57–76.
- [17] A.P. Nolasco and L.P. Castro, Factorization of Wiener-Hopf plus Hankel operators with APW Fourier symbols, *Int. J. Pure Appl. Math.*, **14** (2004), 537–550.
- [18] A.P. Nolasco and L.P. Castro, A criterion for lateral invertibility of matrix Wiener-Hopf plus Hankel operators with good Hausdorff sets, 9 p., *to appear*.

- [19] A.P. Nolasco and L.P. Castro, A Duduchava-Saginashvili's type theory for Wiener-Hopf plus Hankel operators, 16 p., *to appear*.
- [20] H.R. Pousson, Systems of Toeplitz operators on H^2 , *Proc. Amer. Math. Soc.*, **19** (1968), 603–608.
- [21] M. Rabindranathan, On the inversion of Toeplitz operators, *J. Math. Mech.*, **19** (1969/1970), 195–206.
- [22] S. Roch and B. Silbermann, *Algebras of Convolution Operators and their Image in the Calkin Algebra*, Akademie der Wissenschaften der DDR, Karl-Weierstrass-Institut für Mathematik, Berlin (1990).
- [23] F.S. Teixeira, On a class of Hankel operators: Fredholm properties and invertibility, *Integral Equations Operator Theory*, **12** (1989), 592–613.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AVEIRO, 3810-193 AVEIRO, PORTUGAL
E-mail address: `giorgi@mat.ua.pt`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AVEIRO, 3810-193 AVEIRO, PORTUGAL
E-mail address: `lcastro@mat.ua.pt`
URL: `http://www.mat.ua.pt/lcastro/`